


Kiyoshi Tsutsuki http://timetraveler.html.xdomain.jp

рΗ

 $= -\log (H^{+})$

oriba Home page




pH and crop growth (vegetables and root crops)


Low pH tolerance	Crops
strong	Potato, taro
(4.0~5.0)	
A little strong	Sweet potato, white radish, turnip,
(4.5~6.0)	green bean, carrot, cucumber, parsley
A little weak	Tomato, egg plant, cabbage,
(5.5~6.5)	cauliflower, broccoli, celery, green
	peas, melon
weak	Spinach, Onion, leek, burdock,
(6.0~7.0)	asparagus, red pepper, lettus

pH and crop growth (grains and pasture crops)				
Low pH tolerance	Crops			
strong (4.0~5.0)	Rice, tea, tobacco			
A little strong (4.5~6.0)	Wheat, timothy			
A little weak (5.5~6.5)	Adzuki bean, clover, milk vetch			
weak (6.0~7.0)	Sugar beet, barley, rye			

Growth medium pH and the dry matter production of crops: average of 49 crops.



Al toxicity in hydroponic culture



Soil pH and the availability of nutrients

Acidic		Alkaline				
pH High Mid Weak 4.0 4.5 5.0 5.5 6.0		Slight .0 7.	Weak 5 8.0		High 9.0 9.5	рН 10.0
	Nitrog	en (N)				
	Phosph	orus (P)				
	Pottasium	(K)		\rightarrow		
	Sulfur	(S)				
	1	Calciu	m (Ca)	a)		
		u I	Magnesium	(Mg)		
Iron, aluminum (Fe, Al)		Π	1			
Manganese (Mn)		л	1			
Boron (B)						
Copper, zinc (Cu, Zn)	-	N.				-
		1	Molybde	enum (Mo)		

Soil acidity and the growth of crops (1)

A) Damage by hydrogen ion.

B) Damage by active aluminum.

C) Deficiency in calcium and magnesium

D) Deficiency in phosphate.

Binding of phosphate and aluminum.

Soil acidity and the growth of crops (2)

E) Leaching and deficiency of boron.

Decrease in solubility of molybdenum and its deficiency.

 \rightarrow Legumes frequently suffer from molybdenum deficiency.

F) Excess in manganese.

As manganese become more soluble in acidic condition.

Soil acidity and the growth of crops (3)

G) Repression of organic matter decomposition.

By ameliorating the soil acidity, mineralization of organic nitrogen and organic phosphorus increase. $_{\rm o}$

H) Change in microbial flora.

Fungi prefer acidic condition, while bacteria and actinomycete prefer alkaline condition. Soil acidity and the growth of crops (4)

I) Repression of nitrogen fixation.
Optimum pH 6.5~7.5
J) Repression of nitrification.
By liming, the activity of nitrification increases remarkably.


Improvement of soil acidity (1)

• Calcium carbonate (CaCO₃)

Apply 3 times of exchangeable acidity (y_1) . Buffer curve method.

• Gypsum (CaSO₄)

In the sublayer, Al³⁺ is replaced by Ca²⁺. High solubility of gypsum helps the reaction.



Calculation of lime requirement (example) Goal pH 6.5 \rightarrow CaCO₃420 mg / 100g soil = 4.2g / kg = 4.2 kg / tAmount of soil in 1 ha, to the depth 15 cm. $= 100 \text{m} \times 100 \text{m} \times 0.15 \text{m} = 1500 \text{m}^3$ = 1500 Mg = 1500 t (Bulk density = 1) Lime requirement / 1 ha is $4.2 \times 1500 = 6300 \text{ kg}$


Improvement of soil acidity (2)

- Apply phosphates in large amounts, because phosphate is hardly soluble in acidic soil.
- Apply organic matter, to increase the buffer capacity to pH change of the soil.

If the soil is made too alkaline,

Nutrient deficiency occurs. For example, Phosphate, calcium, magnesium, boron, iron, manganese, and zinc.

Buffer curves in various soils and water




Features of 3 types of soils

soil	classification	feature
Uenae	Immature volcanic fallout soil	Coarse particles
Nopporo	Gray terrace soil	Clay rich
Shibetsu	Humic andosoil	Humus rich

Mechanisms of soil acidification


- CO₂ in rain water.
- Aluminum in acidic soil.
- Fertilizer application.
- Acid rain.
- Acid sulfate soil.


Cation holding by soil colloids





 $H^+>Al^{3+}>Ca^{2+}>Mg^{2+}>K^+>Na^+$

cations:









$Al^{3+} + H_2O = Al(OH)^{2+} + H^+$

log K = -4.97(as strong as acetic acid) $log K of acetic acid = -4.76 (25^{\circ}C)$